

A diagrammatic approach to categorification of quantum groups II

Mikhail Khovanov and Aaron D. Lauda

April 15, 2008

Abstract

We categorify one-half of the quantum group associated to an arbitrary Cartan datum.

Cartan data and algebras ${}_{\mathcal{A}}\mathbf{f}$. A *Cartan datum* $(I, \cdot)$ consists of a finite set I and a symmetric bilinear form on $\mathbb{Z}[I]$ taking values in $\mathbb{Z}$, subject to conditions

- $i \cdot i \in \{2, 4, 6, \dots\}$ for any $i \in I$,
- $2\frac{i \cdot j}{i \cdot i} \in \{0, -1, -2, \dots\}$ for any $i \neq j$ in I .

We set $d_{ij} = -2\frac{i \cdot j}{i \cdot i} \in \mathbb{N}$. To a Cartan datum assign a graph Γ with the set of vertices I and an edge between i and j if and only if $i \cdot j \neq 0$.

We recall the definition of the negative half of the quantum group associated to a Cartan datum, following [6]. Let $q_i = q^{\frac{i \cdot i}{2}}$, $[n]_i = q_i^{n-1} + q_i^{n-3} + \dots + q_i^{1-n}$, $[n]_i! = [n]_i[n-1]_i \dots [1]_i$. Let $'\mathbf{f}$ be the free associative algebra over $\mathbb{Q}(q)$ with generators θ_i , $i \in I$ and denote $\theta_i^{(n)} = \theta_i^n / [n]_i!$. We equip $'\mathbf{f}$ with an $\mathbb{N}[I]$ -grading by assigning to θ_i grading i . The tensor square $'\mathbf{f} \otimes '\mathbf{f}$ is an associative algebra with the multiplication

$$(x_1 \otimes x_2)(x'_1 \otimes x'_2) = q^{-|x_2| \cdot |x'_1|} x_1 x'_1 \otimes x_2 x'_2$$

for homogeneous x_1, x_2, x'_1, x'_2 . There is a unique algebra homomorphism $r : '\mathbf{f} \longrightarrow \mathbf{f} \otimes '\mathbf{f}$ given on generators by $r(\theta_i) = \theta_i \otimes 1 + 1 \otimes \theta_i$.

Proposition 1. Algebra $'\mathbf{f}$ carries a unique $\mathbb{Q}(v)$ -bilinear form such that $(1, 1) = 1$ and

- $(\theta_i, \theta_j) = \delta_{i,j}(1 - q_i^2)^{-1}$ for all $i, j \in I$,
- $(x, yy') = (r(x), y \otimes y')$ for $x, y, y' \in '\mathbf{f}$,
- $(xx', y) = (x \otimes x', r(y))$ for $x, x', y \in '\mathbf{f}$.

This bilinear form is symmetric.

The radical $\mathfrak{I}$ of $(,)$ is a two-sided ideal of $'\mathbf{f}$. The bilinear form descends to a non-degenerate bilinear form on the associative $\mathbb{Q}(q)$ -algebra $\mathbf{f} = '\mathbf{f}/\mathfrak{I}$. The $\mathbb{N}[I]$ -grading also descends:

$$\mathbf{f} = \bigoplus_{\nu \in \mathbb{N}[I]} \mathbf{f}_\nu.$$

The quantum version of the Gabber-Kac theorem says the following.

Proposition 2. The ideal $\mathfrak{I}$ is generated by the elements

$$\sum_{a+b=d_{ij}+1} (-1)^a \theta_i^{(a)} \theta_j \theta_i^{(b)}$$

over all $i, j \in I$, $i \neq j$.

Thus, $\mathbf{f}$ is the quotient of $'\mathbf{f}$ by the so-called quantum Serre relations

$$\sum_{a+b=d_{ij}+1} (-1)^a \theta_i^{(a)} \theta_j \theta_i^{(b)} = 0. \quad (1)$$

Denote by $\mathcal{A}\mathbf{f}$ the $\mathbb{Z}[q, q^{-1}]$ -subalgebra of $\mathbf{f}$ generated by the divided powers $\theta_i^{(a)}$, over all $i \in I$ and $a \in \mathbb{N}$.

Algebras $R(\nu)$. As in [4], we consider braid-like planar diagrams, each strand labelled by an element of I , and impose the following relations

$$\begin{array}{c} \text{Diagram: two strands labeled } i \text{ and } j \text{ crossing} \end{array} = \left\{ \begin{array}{ll} 0 & \text{if } i = j, \\ \begin{array}{c} | \quad | \\ i \quad j \end{array} & \text{if } i \cdot j = 0, \\ \begin{array}{c} d_{ij} \bullet \quad | \quad | \\ i \quad j \end{array} + \begin{array}{c} | \quad d_{ji} \bullet \\ i \quad j \end{array} & \text{if } i \cdot j \neq 0. \end{array} \right. \quad (2)$$

$$\begin{array}{c} \bullet \\ \diagup \quad \diagdown \\ i \quad j \end{array} = \begin{array}{c} \diagup \quad \diagdown \\ i \quad j \\ \bullet \end{array} \quad \begin{array}{c} \diagdown \quad \diagup \\ i \quad j \\ \bullet \end{array} = \begin{array}{c} \diagup \quad \diagdown \\ i \quad j \\ \bullet \end{array} \quad \text{for } i \neq j \quad (3)$$

$$\begin{array}{c} \bullet \\ \diagup \quad \diagdown \\ i \quad i \end{array} - \begin{array}{c} \diagdown \quad \diagup \\ i \quad i \\ \bullet \end{array} = \begin{array}{c} | \quad | \\ i \quad i \end{array} \quad (4)$$

$$\begin{array}{c} \diagdown \quad \diagup \\ i \quad i \\ \bullet \end{array} - \begin{array}{c} \diagup \quad \diagdown \\ i \quad i \\ \bullet \end{array} = \begin{array}{c} | \quad | \\ i \quad i \end{array} \quad (5)$$

$$\begin{array}{c} \diagup \quad \diagdown \quad \diagup \\ i \quad j \quad k \end{array} = \begin{array}{c} \diagdown \quad \diagup \quad \diagdown \\ i \quad j \quad k \end{array} \quad \text{unless } i = k \text{ and } i \cdot j \neq 0 \quad (6)$$

$$\begin{array}{c} \diagup \quad \diagdown \quad \diagup \\ i \quad j \quad i \end{array} - \begin{array}{c} \diagdown \quad \diagup \quad \diagdown \\ i \quad j \quad i \end{array} = \sum_{a=0}^{d_{ij}-1} \begin{array}{c} | \quad | \quad | \\ i \quad j \quad i \\ \bullet^a \end{array} \quad \text{if } i \cdot j \neq 0 \quad (7)$$

Example 3. For the Cartan datum $\mathbf{B}_2 = \{i \cdot i = 2, j \cdot j = 4, i \cdot j = -2\}$ we have $d_{ij} = 2$, $d_{ji} = 1$, and the relations involving d_{ij}, d_{ji} are

$$\begin{array}{c} \diagup \quad \diagdown \\ i \quad j \end{array} = \begin{array}{c} | \quad | \\ i \quad j \\ \bullet \end{array} + \begin{array}{c} | \quad | \\ i \quad j \\ \bullet \end{array}$$

$$\begin{array}{c} \diagdown \quad \diagup \\ j \quad i \end{array} = \begin{array}{c} | \quad | \\ j \quad i \\ \bullet \end{array} + \begin{array}{c} | \quad | \\ j \quad i \\ \bullet \end{array}$$

$$\begin{array}{c}
\begin{array}{ccc}
\begin{array}{c} \text{Diagram 1: } i \text{ and } j \text{ cross, } i \text{ on top} \\ i \quad j \quad i \end{array} & - & \begin{array}{c} \text{Diagram 2: } i \text{ and } j \text{ cross, } j \text{ on top} \\ i \quad j \quad i \end{array} \\
= & & \begin{array}{c} \text{Diagram 3: } \bullet \text{ on } i \\ i \quad j \quad i \end{array} + \begin{array}{c} \text{Diagram 4: } \bullet \text{ on } j \\ i \quad j \quad i \end{array}
\end{array} \\
\\
\begin{array}{ccc}
\begin{array}{c} \text{Diagram 5: } j \text{ and } i \text{ cross, } j \text{ on top} \\ j \quad i \quad j \end{array} & - & \begin{array}{c} \text{Diagram 6: } j \text{ and } i \text{ cross, } i \text{ on top} \\ j \quad i \quad j \end{array} \\
= & & \begin{array}{c} \text{Diagram 7: } j \quad i \quad j \text{ (parallel)} \end{array}
\end{array}
\end{array}$$

For each $\nu \in \mathbb{N}[I]$ define the graded ring

$$R(\nu) \stackrel{\text{def}}{=} \bigoplus_{i,j \in \text{Seq}(\nu)} {}_j R(\nu)_i, \quad (8)$$

where ${}_j R(\nu)_i$ is the abelian group of all linear combinations of diagrams with $\text{bot}(D) = \mathbf{i}$ and $\text{top}(D) = \mathbf{j}$ modulo the relations (2)–(7) and $\text{Seq}(\nu)$ is the set of weight ν sequences of elements of I . The multiplication is given by concatenation. Degrees of the generators are

$$\deg \left(\begin{array}{c} | \\ \bullet \\ | \\ i \end{array} \right) = i \cdot i, \quad \deg \left(\begin{array}{cc} & \diagup \\ \diagdown & \\ i & j \end{array} \right) = -i \cdot j. \quad (9)$$

The rest of [4, Section 2.1] generalizes without difficulty to an arbitrary Cartan datum. To define the analogue of the module $\mathcal{Pol}_\nu$ over $R(\nu)$, we choose an orientation of each edge of Γ , then faithfully follow the exposition in Section 2.3 of [4], only changing the action of $\delta_{k,i}$ in the last of the four cases to

$$f \mapsto (x_k(s_k \mathbf{i})^d + x_{k+1}(s_k \mathbf{i})^{d'})(s_k f) \quad \text{if } i_k \longrightarrow i_{k+1},$$

where $d = d_{i_{k+1}i_k}$ and $d' = d_{i_k i_{k+1}}$. Here notation $i_k \longrightarrow i_{k+1}$ means that $i_k \cdot i_{k+1} \neq 0$ and this edge of Γ is oriented from i_k to i_{k+1} . Proposition 2.3 in [4] holds for an arbitrary $(I, \cdot)$. As in [4, Section 2.3], we define ${}_j B_i$, which might depend on minimal presentations of permutations in ${}_j S_i$ and gives a basis in ${}_j R(\nu)_i$. Corollary 2.6 in [4], showing that $\mathcal{Pol}_\nu$ is a faithful graded module over $R(\nu)$, holds for an arbitrary Cartan datum and the properties of $R(\nu)$ established in [4, Section 2.4] generalize without difficulty.

Computations in the nilHecke ring. In this section we slightly enhance the graphical calculus for computations in the nilHecke ring and record several lemmas to be used in the proof of categorified quantum Serre relations below. We use notations from Section 2.2 of [4].

A box with n incoming and n outgoing edges and $\partial(n)$ written inside denotes the longest divided difference ∂_{w_0} , the nonzero product of $\frac{n(n-1)}{2}$ divided differences from $\{\partial_1, \dots, \partial_{n-1}\}$:

The diagram shows a box labeled $\partial(n)$ with n incoming edges at the bottom and n outgoing edges at the top. Below the box is a brace labeled n . This is equal to a braid diagram with n strands. The strands are arranged in a way that they cross each other in a specific pattern, representing the longest divided difference.

When this box is part of a diagram for an element of $R(\nu)$, it denotes the corresponding element of $R(ni) \subset R(\nu)$. A box labelled e_n denotes the idempotent $e_n = x_1^{n-1}x_2^{n-2} \dots x_{n-1}\partial_{w_0}$.

The diagram shows a box labeled e_n with n incoming edges at the bottom and n outgoing edges at the top. This is equal to a box labeled $\partial(n)$ with n incoming edges at the bottom and n outgoing edges at the top. The top $n-1$ strands have dots above them, and the bottom strand has a dot above it.

A box labelled $e_{i,n}$ denotes the corresponding idempotent in $R(\nu)$:


The diagram shows a box labeled $e_{i,n}$ with n incoming edges at the bottom and n outgoing edges at the top. The bottom strand is labeled i . This is equal to a box labeled $\partial(n)$ with n incoming edges at the bottom and n outgoing edges at the top. The top $n-1$ strands have dots above them, and the bottom strand has a dot above it. The bottom strand is labeled i .

Remark 4. Similar-looking diagrams are used in the graphical calculus of Jones-Wenzl projectors, see [3], but the latter has no direct relation to the graphical calculus in our paper.

Lemma 5. We have

The diagram shows a box labeled $\partial(n)$ with n incoming edges at the bottom and n outgoing edges at the top. This is equal to a box labeled $e_{i,n}$ with n incoming edges at the bottom and n outgoing edges at the top. The top $n-1$ strands have dots above them, and the bottom strand has a dot above it.

Proof is by induction on n :



The first equality uses that $x_1 x_2 \dots x_{n-1}$ is central in the nilHecke ring NH_{n-1} , allowing us to move these dots across $\partial(n-1)$. The second equality is the induction hypothesis. $\square$

The lemma implies the following graphical identities:

$$\begin{array}{c} \text{---} \\ \text{---} \\ \boxed{e_n} \\ \text{---} \\ \boxed{e_{n-1}} \\ \text{---} \end{array} = \begin{array}{c} \text{---} \\ \text{---} \\ \boxed{e_n} \\ \text{---} \\ \text{---} \\ \text{---} \end{array} \quad \begin{array}{c} \text{---} \\ \text{---} \\ \boxed{e_n} \\ \text{---} \\ \boxed{e_{n-1}} \\ \text{---} \end{array} = \begin{array}{c} \text{---} \\ \text{---} \\ \boxed{e_n} \\ \text{---} \\ \text{---} \\ \text{---} \end{array} \quad (10)$$

Diagrammatic equation (11) showing the equivalence of two configurations of boxes e_{n-1} and e_n with strands. The left side shows a box e_{n-1} above a box e_n , with four strands entering from the bottom and exiting from the top. The right side shows a box e_{n-1} above a box e_n , with four strands entering from the bottom and exiting from the top, but the strands are rearranged. The equation is labeled (11).

The following also hold:

$$\begin{array}{c} \text{Diagram 1: } e_n \text{ box with } n \text{ top strands and } n \text{ bottom strands. A dot is on the } a\text{-th strand from the left.} \\ = \begin{cases} 0 & \text{if } a < n-1, \\ \text{Diagram 2: } e_n \text{ box with } n \text{ top strands and } n \text{ bottom strands.} & \text{if } a = n-1, \end{cases} \end{array} \quad (12)$$

$$\begin{array}{c} \text{Diagram 3: } e_n \text{ box with } n \text{ top strands and } n \text{ bottom strands. A dot is on the } a\text{-th strand from the right.} \\ = \begin{cases} 0 & \text{if } a < n-1, \\ (-1)^{n-1} \text{Diagram 4: } e_n \text{ box with } n \text{ top strands and } n \text{ bottom strands.} & \text{if } a = n-1. \end{cases} \end{array} \quad (13)$$

For each $i \in I$ the ring $R(mi)$ is isomorphic to the nilHecke ring. The grading of a dot is now $i \cdot i$, while that of a crossing is $-i \cdot i$. For this reason one needs to generalize the grading convention described in [4, Section 2.2] and define ${}_{i,m}P$ to be the right graded projective module $e_{i,m}R(mi)\{-\frac{m(m-1)i \cdot i}{4}\}$, so that the grading starts in the degree $\{-\frac{m(m-1)i \cdot i}{4}\}$. Likewise, $P_{i,m}$ is the left graded projective module $R(mi)\psi(e_{i,m})\{-\frac{m(m-1)i \cdot i}{4}\}$.

The Grothendieck group, bilinear form and projectives. We retain all notations and assumptions from [4], working over a field $\mathbb{k}$, denoting by $K_0(R(\nu))$ the Grothendieck group of the category $R(\nu)\text{-pmod}$ of graded finitely-generated projective left $R(\nu)$ -modules and forming the direct sum

$$R = \bigoplus_{\nu \in \mathbb{N}[I]} R(\nu), \quad K_0(R) = \bigoplus_{\nu \in \mathbb{N}[I]} K_0(R(\nu)).$$

Consider symmetric $\mathbb{Z}[q, q^{-1}]$ -bilinear form

$$(\cdot, \cdot) : K_0(R(\nu)) \times K_0(R(\nu)) \longrightarrow \mathbb{Z}[q^{-1}, q] \cdot (\nu)_q, \quad (14)$$

where

$$(\nu)_q = \text{gdim}(\text{Sym}(\nu)) = \prod_{i \in \Gamma} \left(\prod_{a=1}^{\nu_i} \frac{1}{1 - q^{a \cdot i}} \right), \quad (15)$$

and

$$([P], [Q]) = \text{gdim}_{\mathbb{k}}(P^\psi \otimes_{R(\nu)} Q). \quad (16)$$

The character $\text{ch}(M)$ of an $R(\nu)$ -module M , the divided power sequences $\text{Seqd}(\nu)$, and idempotents $1_{\mathbf{i}}$ for $\mathbf{i} = i_1^{(n_1)} \dots i_r^{(n_r)} \in \text{Seqd}(\nu)$ are defined as in [4, Section 2.5]. Let $\mathbf{i}! = [n_1]_{i_1}! \dots [n_r]_{i_r}!$ and

$$\langle \mathbf{i} \rangle = \sum_{k=1}^r \frac{n_k(n_k - 1)}{2} \cdot \frac{i_k \cdot i_k}{2}.$$

Define graded left, respectively right, projective module

$$P_{\mathbf{i}} = R(\nu)\psi(1_{\mathbf{i}})\{-\langle \mathbf{i} \rangle\}, \quad {}_{\mathbf{i}}P = 1_{\mathbf{i}}R(\nu)\{-\langle \mathbf{i} \rangle\}.$$

Quantum Serre relations. Let

$$\alpha_{a,b}^+(i, j) = \begin{array}{c} \begin{array}{cc} \overbrace{\quad\quad\quad}^{a+1} & \overbrace{\quad\quad\quad}^{b-1} \\ \begin{array}{|c|c|c|c|} \hline e_{i,a+1} \\ \hline \end{array} & \begin{array}{|c|c|} \hline e_{i,b-1} \\ \hline \end{array} \\ \begin{array}{|c|c|c|c|} \hline i \\ \hline \end{array} & \begin{array}{|c|c|c|c|} \hline i \\ \hline \end{array} \end{array} \end{array}, \quad (17)$$

and also write $\alpha_{a,b}^+$ when i and j are fixed. To prove the categorified quantum Serre relations, we assume that $a + b = d + 1$, where $d = d_{ij}$. The element $\alpha_{a,b}^+$ belongs to

$${}_{i^{a+1}ji^{b-1}}R(j + (a+b)i)_{i^a j i^b}.$$

By adding vertical lines on the left and on the right of the diagram, $\alpha_{a,b}^+$ can be viewed, more generally, as an element of

$${}_{\mathbf{i}' i^{a+1} j i^{b-1} \mathbf{i}''}R(\nu)_{\mathbf{i}' i^a j i^b \mathbf{i}''}$$

for any sequences $\mathbf{i}'$, $\mathbf{i}''$ and the corresponding ν . We can replace sequences $\mathbf{i}'$ and $\mathbf{i}''$ by dots to simplify notation.

Left multiplication by $\alpha_{a,b}^+$ is a homomorphism of projective modules

$${}_{\dots i^a j i^b \dots}P \longrightarrow {}_{\dots i^{a+1} j i^{b-1} \dots}P. \quad (18)$$

The top part of the diagram of $\alpha_{a,b}^+$ contains idempotents $e_{i,a+1}$ and $e_{i,b-1}$. Therefore, $\alpha_{a,b}^+$ induces a homomorphism of projective modules

$$\dots i^{(a)} j i^{(b)} \dots P \longrightarrow \dots i^{(a+1)} j i^{(b-1)} \dots P, \quad (19)$$

denoted $\alpha_{(a,b)}^+$ and given by the composition

$$\dots i^{(a)} j i^{(b)} \dots P \hookrightarrow \dots i^a j i^b \dots P \xrightarrow{\alpha_{a,b}^+} \dots i^{(a+1)} j i^{(b-1)} \dots P. \quad (20)$$

It is easy to check that $\alpha_{(a,b)}^+$ is a grading-preserving homomorphism. Likewise, let

$$\alpha_{a,b}^-(i, j) = \begin{array}{c} \begin{array}{c} \overbrace{\quad\quad\quad}^{a-1} \\ \begin{array}{|c|c|c|c|c|} \hline e_{i,a-1} \\ \hline \end{array} \\ \underbrace{\quad\quad\quad}_a \end{array} \quad \begin{array}{c} \overbrace{\quad\quad\quad}^{b+1} \\ \begin{array}{|c|c|c|c|c|} \hline e_{i,b+1} \\ \hline \end{array} \\ \underbrace{\quad\quad\quad}_b \end{array} \end{array} \quad (21)$$

and write $\alpha_{a,b}^-$ instead of $\alpha_{a,b}^-(i, j)$ when i and j are fixed. This element of

$$\dots i^{a-1} j i^{b-1} \dots R(\nu) \dots i^a j i^b \dots \quad (22)$$

gives rise to a grading-preserving homomorphism of projectives

$$\alpha_{(a,b)}^- : \dots i^{(a)} j i^{(b)} \dots P \longrightarrow \dots i^{(a-1)} j i^{(b+1)} \dots P. \quad (23)$$

For the next few pages, denote $\dots i^{(a)} j i^{(b)} \dots P$ by $_{(a,b)}P$ (recall that $a + b = d + 1$, $d = d_{ij}$). We have a diagram of projective modules and grading-preserving homomorphisms

$$\begin{array}{ccccc} \xrightarrow{\quad} & & \xrightarrow{\alpha_{(a-1,b+1)}^+} & & \xrightarrow{\alpha_{(a,b)}^+} \\ \xleftarrow{\quad} & _{(a-1,b+1)}P & \xleftarrow{\alpha_{(a,b)}^-} & _{(a,b)}P & \xleftarrow{\alpha_{(a+1,b-1)}^-} \\ & & & & \end{array} \quad (24)$$

terminating on the left at

$$\begin{array}{ccc} & \xrightarrow{\alpha_{(0,d+1)}^+} & \\ (0,d+1)P & \xleftarrow{\alpha_{(1,d)}^-} & (1,d)P \xleftarrow{\quad} \dots \end{array} \quad (25)$$

and on the right at

$$\cdots \begin{array}{c} \xrightarrow{\quad} \\ \xleftarrow{\quad} \end{array} (d,1)P \begin{array}{c} \xrightarrow{\alpha_{(d,1)}^+} \\ \xleftarrow{\alpha_{(d+1,0)}^-} \end{array} (d+1,0)P. \quad (26)$$

Relations (10), (11) imply

$$\alpha_{a+1,b-1}^- \alpha_{a,b}^+ =$$

$$\stackrel{(10)}{=} \quad \stackrel{(11)}{=}$$

Furthermore,

$$\alpha_{a-1,b+1}^+ \alpha_{a,b}^- =$$

$$=$$

Therefore,

$$\alpha_{a-1,b+1}^+ \alpha_{a,b}^- - \alpha_{a+1,b-1}^- \alpha_{a,b}^+ = (-1)^{a-1} e_{i,a} \otimes 1_j \otimes e_{i,b},$$

as elements of $R(\nu)$, see below:

$$\begin{aligned} \alpha_{a-1,b+1}^+ \alpha_{a,b}^- - \alpha_{a+1,b-1}^- \alpha_{a,b}^+ &= \begin{array}{c} \text{Diagram 1} \end{array} - \begin{array}{c} \text{Diagram 2} \end{array} \\ &\stackrel{(7)}{=} \sum_{c=0}^{d-1} \begin{array}{c} \text{Diagram 3} \end{array} \stackrel{(12),(13)}{=} \begin{array}{c} \text{Diagram 4} \end{array} \\ &\stackrel{(12),(13)}{=} (-1)^{a-1} \begin{array}{c} \text{Diagram 5} \end{array}, \end{aligned}$$

The diagrams are as follows:

- Diagram 1:** Two boxes labeled $e_{i,a}$ and $e_{i,b}$. The $e_{i,a}$ box has 4 inputs from below labeled i, i, i, i and 4 outputs from above. The $e_{i,b}$ box has 3 inputs from below labeled i, i, i and 3 outputs from above. A vertical line separates the two boxes. The j input is connected to the $e_{i,b}$ box.
- Diagram 2:** Similar to Diagram 1, but the j input is connected to the $e_{i,a}$ box.
- Diagram 3:** Two boxes labeled $e_{i,a}$ and $e_{i,b}$. The $e_{i,a}$ box has 4 inputs from below labeled i, i, i, i and 4 outputs from above. The $e_{i,b}$ box has 3 inputs from below labeled i, i, i and 3 outputs from above. A vertical line separates the two boxes. The j input is connected to the $e_{i,b}$ box. A dot labeled c is on the i input of the $e_{i,b}$ box. A dot labeled $d-1-c$ is on the i input of the $e_{i,a}$ box.
- Diagram 4:** Similar to Diagram 3, but the dot labeled $a-1$ is on the i input of the $e_{i,a}$ box and the dot labeled $b-1$ is on the i input of the $e_{i,b}$ box.
- Diagram 5:** Two boxes labeled $e_{i,a}$ and $e_{i,b}$. The $e_{i,a}$ box has 4 inputs from below labeled i, i, i, i and 4 outputs from above. The $e_{i,b}$ box has 3 inputs from below labeled i, i, i and 3 outputs from above. A vertical line separates the two boxes. The j input is connected to the $e_{i,b}$ box.

where $d = a + b - 1$. Consequently,

$$\alpha_{(a-1,b+1)}^+ \alpha_{(a,b)}^- - \alpha_{(a+1,b-1)}^- \alpha_{(a,b)}^+ = (-1)^{a-1} \cdot \text{Id},$$

as endomorphisms of the projective module ${}_{(a,b)}P$, since $e_{i,a} \otimes 1_j \otimes e_{i,b}$ acts by the identity on ${}_{(a,b)}P$.

Likewise,

$$\begin{aligned}
 \alpha_{1,d}^- \alpha_{0,d+1}^+ &= \text{Diagram 1} \stackrel{(3)}{=} \text{Diagram 2} + \text{Diagram 3} \\
 &\stackrel{(12)}{=} \text{Diagram 4} \stackrel{(12)}{=} \text{Diagram 5} = 1_j \otimes e_{i,d+1},
 \end{aligned}$$

The diagrams are as follows:

- Diagram 1:** A box labeled $e_{i,d+1}$ is connected to a box labeled $e_{i,d}$. The input lines from the bottom are labeled j, i, i, i, i . The output lines from the top are labeled j, i, i, i, i .
- Diagram 2:** A box labeled $e_{i,d+1}$ with a dot labeled d' on the input line j . The input lines from the bottom are labeled j, i, i, i, i . The output lines from the top are labeled j, i, i, i, i .
- Diagram 3:** A box labeled $e_{i,d+1}$ with a dot labeled d on the input line j . The input lines from the bottom are labeled j, i, i, i, i . The output lines from the top are labeled j, i, i, i, i .
- Diagram 4:** A box labeled $e_{i,d+1}$ with a dot labeled d on the input line j . The input lines from the bottom are labeled j, i, i, i, i . The output lines from the top are labeled j, i, i, i, i .
- Diagram 5:** A box labeled $e_{i,d+1}$ with input lines from the bottom labeled j, i, i, i, i and output lines from the top labeled j, i, i, i, i .

where $d' = d_{ji}$, and

$$\alpha_{(1,d)}^- \alpha_{(0,d+1)}^+ = \text{Id},$$

as endomorphisms of ${}_{(0,d+1)}P$. A similar computation shows that

$$\alpha_{d,1}^+ \alpha_{d+1,0}^- = (-1)^d e_{i,d+1} \otimes 1_j,$$

as elements of $R(\nu)$, and

$$\alpha_{(d,1)}^+ \alpha_{(d+1,0)}^- = (-1)^d \cdot \text{Id},$$

as endomorphisms of ${}_{(d+1,0)}P$.

Proposition 6. For each $i, j \in I, i \neq j$ there are isomorphisms of graded right projective modules

$$\bigoplus_{a=0}^{\lfloor \frac{d+1}{2} \rfloor} \dots i^{(2a)} j i^{(d+1-2a)} \dots P \cong \bigoplus_{a=0}^{\lfloor \frac{d}{2} \rfloor} \dots i^{(2a+1)} j i^{(d-2a)} \dots P.$$

Proof. When $i \cdot j = 0$ the isomorphism reads

$$\dots j i \dots P \cong \dots i j \dots P$$

and is given by left multiplication by the ij intersection. When $i \cdot j < 0$, earlier computations show that the maps

$$\bigoplus_{a=0}^{\lfloor \frac{d+1}{2} \rfloor} (2a, d+1-2a)P \begin{matrix} \xrightarrow{\alpha'} \\ \xleftarrow{\alpha''} \end{matrix} \bigoplus_{a=0}^{\lfloor \frac{d}{2} \rfloor} (2a+1, d-2a)P$$

given by

$$\begin{aligned} \alpha' &= \sum_{a=0}^{\lfloor \frac{d}{2} \rfloor} \alpha_{(2a, d+1-2a)}^+ + \sum_{a=0}^{\lfloor \frac{d+1}{2} \rfloor} \alpha_{(2a, d+1-2a)}^-, \\ \alpha'' &= \sum_{a=0}^{\lfloor \frac{d+1}{2} \rfloor} \alpha_{(2a+1, d-2a)}^- - \sum_{a=0}^{\lfloor \frac{d}{2} \rfloor} \alpha_{(2a+1, d-2a)}^+, \end{aligned}$$

are mutually-inverse isomorphisms, implying the proposition. Maps α', α'' together are given by summing over all arrows in the diagram (24), with every fourth arrow appearing with the minus sign. $\square$

Corollary 7. For each $i, j \in I, i \neq j$ there are isomorphisms of graded left projective modules

$$\bigoplus_{a=0}^{\lfloor \frac{d+1}{2} \rfloor} P_{\dots i^{(2a)} j i^{(d+1-2a)} \dots} \cong \bigoplus_{a=0}^{\lfloor \frac{d}{2} \rfloor} P_{\dots i^{(2a+1)} j i^{(d-2a)} \dots}$$

Proposition 6 and Corollary 7 generalize Proposition 2.13 in [4] and can be considered a categorification of the quantum Serre relations. Corollaries 2.14 and 2.15 of [4], establishing quantum Serre relations for the characters of any $M \in R(\nu)\text{-mod}$, generalize to an arbitrary Cartan datum in the same way.

Grothendieck group as the quantum group. Induction and restriction functors for inclusions $R(\nu) \otimes R(\nu') \subset R(\nu + \nu')$ turn $K_0(R)$ into a twisted bialgebra, and all results of [4, Section 2.6] remain valid for an arbitrary Cartan datum. As in [4, Section 3.1] we define a homomorphism of twisted bialgebras

$$\gamma : \mathcal{A}\mathbf{f} \longrightarrow K_0(R)$$

which takes the product of divided powers $\theta_{\mathbf{i}} = \theta_{i_1}^{(n_1)} \dots \theta_{i_r}^{(n_r)}$ to $[P_{\mathbf{i}}]$, where $\mathbf{i} = i_{i_1}^{(n_1)} \dots i_{i_r}^{(n_r)}$. Homomorphism γ intertwines the bilinear forms on $\mathcal{A}\mathbf{f}$ and $K_0(R)$,

$$(x, y) = (\gamma(x), \gamma(y)), \quad x, y \in \mathcal{A}\mathbf{f}.$$

Due to the quantum Gabber-Kac theorem, this homomorphism is injective. Surjectivity of γ follows from the arguments identical to those given in [4, Section 3.2], which, in turn, were adopted from [5, Section 5]. Alternatively, the arguments could be adopted from [2] and [9]; we settled on using a single source. We obtain

Theorem 8. $\gamma : \mathcal{A}\mathbf{f} \longrightarrow K_0(R)$ is an isomorphism of $\mathbb{N}[I]$ -graded twisted bialgebras.

This theorem holds without any restrictions on the Cartan datum and on the ground field $\mathbb{k}$ over which $R(\nu)$ is defined. All other results and observations of Sections 3.2 and 3.3 of [4] extend to the general case as well. The cyclotomic quotients of $R(\nu)$ described in [4, Section 3.4] generalize to an arbitrary Cartan datum.

It would be interesting to relate our construction to Lusztig's geometric realization of U^- in the non-simply laced case [6] and to Brundan-Kleshchev's categorification [1], [5] of $U_{q=1}^-$ in the affine Dynkin case $A_n^{(2)}$.

A multi-grading. For every pair (i, j) of vertices of Γ , algebras $R(\nu)$ can be equipped with an additional grading, by assigning degrees -1 and 1 to the ij and ji crossings, respectively,

$$\deg \left(\begin{array}{c} \diagup \quad \diagdown \\ i \quad j \end{array} \right) = -1, \quad \deg \left(\begin{array}{c} \diagdown \quad \diagup \\ j \quad i \end{array} \right) = 1,$$

and degree 0 to all other diagrammatic generators of $R(\nu)$. These gradings are independent, and together with the principal grading, introduced above, make $R(\nu)$ into a multi-graded ring (with $\frac{n(n-1)}{2} + 1$ independent gradings where $n = |\text{Supp}(\nu)|$). The direct sum of the categories of multi-graded finitely-generated projective left $R(\nu)$ -modules, over all $\nu \in \mathbb{N}[I]$, categorifies a multi-parameter deformation [7], [8] of the quantum universal enveloping algebra U^- , the quotient of the free associative algebra on θ_i , $i \in I$, by the relations

$$\sum_{a+b=d_{ij}+1} (-1)^a q_{ij}^a \theta_i^{(a)} \theta_j \theta_i^{(b)} = 0, \quad (27)$$

where q_{ij} are formal variables subject to conditions $q_{ij}q_{ji} = 1$.

Modifications in the simply-laced case. This section explains how to deform algebras $R(\nu)$ in the simply-laced case so that the main results of [4] will hold for the modified algebras. These deformations can be nontrivial only when the graph has cycles. As in [4], we start with an unoriented graph Γ without loops and multiple edges. Next, fix an orientation of each edge of Γ , work over a base field $\mathbb{k}$, and, for each oriented edge $i \longrightarrow j$, choose two invertible elements τ_{ij} and τ_{ji} in $\mathbb{k}$. Denote such a datum $\{\text{orientations, invertible elements}\}$ by τ .

For each $\nu \in \mathbb{N}[I]$ consider $\mathbb{k}$ -vector space $\mathcal{Pol}_\nu$ defined as in [4]. This space is the sum of polynomial rings in $|\nu|$ variables, over all sequences in $\text{Seq}(\nu)$. Define $R_\tau(\nu)$ to be the endomorphism algebra of $\mathcal{Pol}_\nu$ generated by the endomorphisms $1_i, x_{k,i}, \delta_{k,i}$, over all possible k and $\mathbf{i}$, with the action as in [4, Section 2.3], with the only difference being the action of $\delta_{k,i}$ in the last of the four cases:

$$f \mapsto (\tau_{i_k i_{k+1}} x_{k+1}(s_k \mathbf{i}) - \tau_{i_{k+1} i_k} x_k(s_k \mathbf{i}))(s_k f) \quad \text{if } i_k \longrightarrow i_{k+1},$$

instead of

$$f \mapsto (x_k(s_k \mathbf{i}) + x_{k+1}(s_k \mathbf{i}))(s_k f) \quad \text{if } i_k \longrightarrow i_{k+1}.$$

The algebra $R_\tau(\nu)$ has a diagrammatic description similar to that of $R(\nu)$, with the following defining relations

$$\begin{array}{c} \text{Diagram of two crossing lines labeled } i \text{ and } j \end{array} = \left\{ \begin{array}{ll} 0 & \text{if } i = j, \\ \begin{array}{c} | \quad | \\ i \quad j \end{array} & \text{if } i \cdot j = 0, \\ \tau_{ij} \begin{array}{c} | \\ \bullet \\ i \end{array} \begin{array}{c} | \\ j \end{array} - \tau_{ji} \begin{array}{c} | \\ i \end{array} \begin{array}{c} | \\ \bullet \\ j \end{array} & \text{if } i \longrightarrow j, \\ \tau_{ji} \begin{array}{c} | \\ i \end{array} \begin{array}{c} | \\ \bullet \\ j \end{array} - \tau_{ij} \begin{array}{c} | \\ \bullet \\ i \end{array} \begin{array}{c} | \\ j \end{array} & \text{if } i \longleftarrow j, \end{array} \right. \quad (28)$$

$$\begin{array}{c} \bullet \\ \diagup \quad \diagdown \\ i \quad j \end{array} = \begin{array}{c} \diagup \quad \diagdown \\ i \quad j \\ \bullet \end{array} \quad \begin{array}{c} \diagdown \quad \diagup \\ i \quad j \\ \bullet \end{array} = \begin{array}{c} \diagdown \quad \diagup \\ i \quad j \\ \bullet \end{array} \quad \text{for } i \neq j \quad (29)$$

$$\begin{array}{c} \bullet \\ \diagup \quad \diagdown \\ i \quad i \end{array} - \begin{array}{c} \diagdown \quad \diagup \\ i \quad i \\ \bullet \end{array} = \begin{array}{c} | \quad | \\ i \quad i \end{array} \quad (30)$$

$$\begin{array}{c} \diagdown \quad \diagup \\ i \quad i \\ \bullet \end{array} - \begin{array}{c} \diagup \quad \diagdown \\ i \quad i \\ \bullet \end{array} = \begin{array}{c} | \quad | \\ i \quad i \end{array} \quad (31)$$

$$\begin{array}{c} \diagup \quad \diagdown \quad \diagup \\ i \quad j \quad k \end{array} = \begin{array}{c} \diagup \quad \diagdown \quad \diagup \\ i \quad j \quad k \end{array} \quad \text{unless } i = k \text{ and } i \cdot j = -1 \quad (32)$$

$$\begin{array}{c} \diagup \quad \diagdown \quad \diagup \\ i \quad j \quad i \end{array} - \begin{array}{c} \diagup \quad \diagdown \quad \diagup \\ i \quad j \quad i \end{array} = \tau_{ij} \begin{array}{c} | \quad | \quad | \\ i \quad j \quad i \end{array} \quad \text{if } i \longrightarrow j \quad (33)$$

$$\begin{array}{c} \diagdown \quad \diagup \quad \diagdown \\ i \quad j \quad i \end{array} - \begin{array}{c} \diagdown \quad \diagup \quad \diagdown \\ i \quad j \quad i \end{array} = -\tau_{ij} \begin{array}{c} | \quad | \quad | \\ i \quad j \quad i \end{array} \quad \text{if } i \longleftarrow j \quad (34)$$

Reverse the orientation of a single edge $i - j$ and change τ_{ij} to $-\tau_{ij}$ and τ_{ji} to $-\tau_{ji}$. Denote the new datum by τ' . Algebras $R_\tau(\nu)$ and $R_{\tau'}(\nu)$ are isomorphic via a map which is the identity on diagrams. This way, the study of $R_\tau(\nu)$ reduces to the case of any preferred orientation of Γ . Rescaling one of the two possible types of the ij crossing by $\lambda \in \mathbb{k}$ changes τ_{ij} to $\lambda\tau_{ij}$ and τ_{ji} to $\lambda\tau_{ji}$ while keeping the rest of the data fixed. We see that $R_\tau(\nu)$ depends only on products $\tau_{ij}\tau_{ji}^{-1}$, over all edges of Γ , via non-canonical isomorphisms. Rescalings of ii crossings and dots further reduce the number

of parameters to the rank of the first homology group of Γ . When graph Γ is a forest (has no cycles), algebras $R_\tau(\nu)$ are all isomorphic to $R(\nu)$ via rescaling of generators. When Γ has a single cycle, rescaling of generators reduces this family of algebras to a one-parameter family, with the parameter taking values in $\mathbb{k}^*$. It is likely that $R_\tau(\nu)$ has a description via equivariant convolution algebras in Lusztig's geometrization [6] of U^- when all $\tau_{ij} = 1$ (compare with Conjecture 1.2 in [4]).

Form

$$R_\tau = \bigoplus_{\nu \in \mathbb{N}[I]} R_\tau(\nu).$$

The Grothendieck group $K_0(R_\tau)$ of the category of finitely-generated graded left projective modules can be naturally identified with the integral version $\mathcal{A}\mathbf{f}$ of U^- . All other essential constructions and results of [4] generalize from $R(\nu)$ to algebras $R_\tau(\nu)$ in a straightforward fashion.

Modifications in the general case. Rings $R(\nu)$ associated to an arbitrary Cartan datum admit similar modifications that depend on choosing an orientation of Γ and invertible elements τ_{ij}, τ_{ji} of the ground field $\mathbb{k}$ for each oriented edge $i \longrightarrow j$. The key point is the change in the definition of the endomorphism algebra, making $\delta_{k,i}$ act by

$$f \mapsto (\tau_{i_k i_{k+1}} x_{k+1} (s_k \mathbf{i})^{d'} - \tau_{i_{k+1} i_k} x_k (s_k \mathbf{i})^d) (s_k f) \quad \text{if } i_k \longrightarrow i_{k+1}$$

in the last of the four cases, with $d = d_{i_{k+1} i_k}$ and $d' = d_{i_k i_{k+1}}$. Our proof of categorified quantum Serre relations for $R(\nu)$ requires only minor changes in the general case of $R_\tau(\nu)$. Everything else generalizes as well.

Acknowledgments. M.K. was fully supported by the IAS and the NSF grants DMS-0635607 and DMS-0706924 while working on this paper.

References

- [1] J. Brundan and A. Kleshchev. Hecke-Clifford superalgebras, crystals of type $A_{2l}^{(2)}$ and modular branching rules for $\widehat{S}_n$. *Represent. Theory*, 5:317–403 (electronic), 2001, math.RT/0103060.
- [2] I. Grojnowski and M. Vazirani. Strong multiplicity one theorems for affine Hecke algebras of type A. *Transform. Groups*, 6(2):143–155, 2001.

- [3] L. Kauffman and S. Lins. *Temperley-Lieb recoupling theory and invariants of 3-manifolds*, volume 134 of *Annals of Mathematics Studies*. Princeton University Press, 1994.
- [4] M. Khovanov and A. Lauda. A diagrammatic approach to categorification of quantum groups I, 2008, math.QA/0803.4121.
- [5] A. Kleshchev. *Linear and projective representations of symmetric groups*, volume 163 of *Cambridge Tracts in Mathematics*. Cambridge U. Press, 2005.
- [6] G. Lusztig. *Introduction to quantum groups*, volume 110 of *Progress in Mathematics*. Birkhäuser Boston Inc., Boston, MA, 1993.
- [7] M. Okado and H. Yamane. R -matrices with gauge parameters and multi-parameter quantized enveloping algebras. In *Special functions (Okayama, 1990)*, ICM-90 Satell. Conf. Proc., pages 289–293. Springer, Tokyo, 1991.
- [8] N. Reshetikhin. Multiparameter quantum groups and twisted quasitriangular Hopf algebras. *Lett. Math. Phys.*, 20(4):331–335, 1990.
- [9] M. Vazirani. Irreducible modules over the affine Hecke algebra: a strong multiplicity one result. Ph.D. thesis, UC Berkeley, 1999, math.RT/0107052.

M.K.: *School of Mathematics, Institute for Advanced Study, Princeton, NJ 08540,*
and Department of Mathematics, Columbia University, New York, NY 10027
email: khovanov@math.columbia.edu

A.L: *Department of Mathematics, Columbia University, New York, NY 10027*
email: lauda@math.columbia.edu